

## Phytopharmacology: Nature's Defense Against Oxidative Stress

Jyoti Kanwa<sup>1\*</sup>, Amrita Suryavanshi<sup>1</sup>

<sup>1\*</sup>Government P.G. College, Safidon, Haryana

<sup>1</sup>Medicinal Plants Research Laboratory, Department of Botany, University of Delhi

### Abstract

Secondary metabolites, a diverse group of organic compounds found in various organisms, play a pivotal role in mitigating oxidative stress—a condition characterized by the imbalance between the production of reactive oxygen species (ROS) and the body's antioxidant defenses. These compounds, not essential for an organism's basic functions but critical for ecological interactions and defense mechanisms, exhibit an astonishing chemical diversity that encompasses alkaloids, terpenoids, phenolic compounds, polyketides, glycosides, lignans, and saponins. Secondary metabolites act as antioxidants through multiple mechanisms, including scavenging ROS, metal chelation, enhancing enzymatic antioxidant systems, and regenerating other antioxidants. However, challenges such as bioavailability, standardization, environmental concerns, and regulatory issues need to be addressed to harness the full potential of secondary metabolites. Despite these challenges, secondary metabolites remain a fascinating area of research with the potential to unlock innovative solutions for human health. This comprehensive review examines the mechanisms underlying oxidative stress, explores the antioxidant properties of phytochemicals, and discusses the potential of phytopharmacology in preventing and managing oxidative stress-related conditions.

**Keywords:** Metabolites, organic, antioxidant, compound, phytochemicals

### 1. Introduction

Secondary metabolites are a diverse group of organic compounds found in various organisms, including plants, microbes, and fungi. These compounds, often referred to as natural products, are not directly involved in the growth, development, or reproduction of the organism but play critical roles in ecological interactions and defense mechanisms (Badyal et al., 2020). Phytopharmacology, the use of plant-based remedies for therapeutic purposes, has emerged as a promising avenue in the battle against oxidative stress. In this article, we delve into the science behind oxidative stress and explore how phytopharmacology offers natural solutions.

#### 1.1. Oxidative Stress: The Cellular Culprit

Oxygen is essential for life, but it can also be a double-edged sword. While the body's cells utilize oxygen to produce energy, this process generates ROS (reactive oxygen species) as byproducts. In normal circumstances, the body maintains a delicate balance between ROS production and antioxidant defense mechanisms. However, this balance can be disrupted, when ROS overwhelm the body's antioxidant capacity leading to oxidative stress.

This excess of ROS can damage vital biomolecules such as DNA, proteins, and lipids, is associated with various health issues, including aging, neurodegenerative diseases, cardiovascular diseases, and cancer (Bhatti et al., 2017). To combat oxidative stress and its detrimental effects, organisms have developed intricate defense mechanisms, including the production of antioxidant secondary metabolites.

#### Sources of ROS

ROS can originate from both endogenous and exogenous sources (Banik and Bhattacharjee, 2020):

##### a. Endogenous Sources:

ROS are produced as byproducts of mitochondrial respiration, enzymatic reactions (e.g., NADPH oxidase), and the peroxisomal metabolism of fatty acids.

**Mitochondrial Respiration:** During the electron transport chain (ETC) in mitochondria, electrons can leak from the respiratory complexes and react with oxygen molecules, leading to the formation of superoxide anion radicals ( $O_2^{\bullet-}$ ). Superoxide is a primary ROS generated within the mitochondria (Kowalczyk et al., 2021). When not efficiently detoxified, it can cause damage to mitochondrial DNA, proteins, and lipids.

**NADPH Oxidase (NOX) Enzymes:** NADPH oxidases are a family of enzymes that produce ROS as part of the immune system's defense mechanism. These enzymes are responsible for generating superoxide and other ROS to help destroy invading pathogens, such as bacteria and viruses (Whitehead et al., 2010).

**Peroxisomes:** Peroxisomes are cellular organelles involved in various metabolic processes, including the breakdown of fatty acids. During these processes, hydrogen peroxide ( $H_2O_2$ ) is produced as a byproduct.  $H_2O_2$  is a relatively stable

ROS, but its accumulation can still lead to oxidative stress if not efficiently neutralized by cellular antioxidants (Pizzino et al., 2017).

### **b. Exogenous Sources:**

External factors, such as ionizing radiation, ultraviolet (UV) radiation, pollution, and certain drugs, can also lead to ROS formation.

**Ultraviolet (UV) Radiation:** Ultraviolet radiation from the sun can directly generate ROS in skin cells. UV radiation can excite electrons in cellular molecules, leading to the formation of ROS such as singlet oxygen and hydroxyl radicals (Ryter and Tyrrell, 1998). Chronic exposure to UV radiation is a major risk factor for skin aging and skin cancers due to its ROS-inducing effects.

### **Consequences of Oxidative Stress**

Excessive ROS can damage lipids, proteins, and DNA, leading to various pathological conditions, including:

**a. Lipid Peroxidation:** Oxidative stress can initiate lipid peroxidation, a chain reaction in which ROS attack and damage the lipids in cell membranes (Mohamed and Hunter, 1986). This process results in the production of lipid peroxidation products, such as malondialdehyde (MDA) and 4-hydroxynonenal (4-HNE), which can further damage cellular structures. Lipid peroxidation is associated with several chronic diseases, including atherosclerosis, neurodegenerative disorders, and liver diseases.

**b. Protein Oxidation:** Proteins are essential for various cellular functions, and oxidative stress can lead to protein oxidation. ROS can modify amino acid residues in proteins, causing changes in their structure and function (McCord, 2000). This can result in the loss of enzymatic activity, misfolding, and aggregation of proteins. Protein oxidation is implicated in age-related diseases, including Alzheimer's disease and Parkinson's disease.

**c. DNA Damage:** Oxidative stress-induced DNA damage can lead to mutations and genomic instability. DNA lesions caused by ROS can result in base modifications, single-strand breaks, and double-strand breaks. Accumulated DNA damage over time is a driving force behind aging and can increase the risk of cancer development (Martins et al., 2021). Cells have repair mechanisms to counteract DNA damage, but when oxidative stress overwhelms these repair systems, mutations can accumulate.

**d. Cellular Dysfunction:** One of the most immediate and direct consequences of oxidative stress is cellular damage. ROS, including superoxide radicals ( $O_2^{\bullet-}$ ), hydrogen peroxide ( $H_2O_2$ ), and hydroxyl radicals ( $OH^{\bullet}$ ), are highly reactive molecules that can interact with cellular components such as DNA, proteins, and lipids. These interactions can lead to oxidative modifications, causing structural and functional damage to biomolecules, including neurodegenerative disorders (e.g., Alzheimer's and Parkinson's), cardiovascular diseases, cancer, and aging-related conditions (Leenders et al., 2021).

## **1.2. Secondary Metabolites: Nature's Chemical Arsenal**

### **1.2.1. Plant-Based Antioxidants:**

**a. Phenolic Acids:** These compounds are characterized by their aromatic rings and hydroxyl groups such as flavonoids and tannins etc. Phenolic acids are found in various plant-based foods, including whole grains, fruits, and vegetables. They help protect cells from oxidative damage and inflammation (Maleki et al., 2017).

**b. Flavonoids:** Flavonoids are a diverse group of polyphenolic compounds found in fruits, vegetables, tea, and red wine. They are known for their potent antioxidant properties and have been linked to a reduced risk of chronic diseases.

**c. Carotenoids:** Carotenoids, such as beta-carotene, lutein, and zeaxanthin, are pigments found in colorful fruits and vegetables. They act as antioxidants and are essential for body health and immunity (Gammone et al., 2015).

**d. Terpenoids:** Terpenoids are derived from isoprene units, these compounds are abundant in plant essential oils and contribute to various plant defense mechanisms. Some terpenoids, like resveratrol found in grapes, have antioxidant properties (Ozcan et al., 2015). Resveratrol has gained attention for its potential cardiovascular benefits.

### **1.3. Antioxidant Defense Mechanism**

Secondary metabolites can act as antioxidants through various mechanisms:

**1.3.1. Scavenging ROS:** Many secondary metabolites (polyphenols, carotenoids, glucosinolates, alkaloids, terpenoids) possess the ability to directly neutralize ROS by donating electrons or hydrogen atoms, thereby preventing ROS-induced damage to cellular components (Banik and Bhattacharjee, 2020).

**1.3.2. Metal Chelation:** Some secondary metabolites (siderophores, phytochelators, tannins, microbial metallothioneins) can bind to transition metal ions, such as iron and copper, which are involved in ROS generation (Puthur et al., 2021).

**1.3.3. Enhancing Enzymatic Antioxidants:** Certain secondary metabolites can upregulate the activity of endogenous antioxidant enzymes, enhancing the cell's defense against oxidative stress (Gorni et al., 2020).

**1.3.4. Regeneration of Other Antioxidants:** Secondary metabolites like vitamins C and E can regenerate other antioxidants, such as glutathione, ensuring a continuous supply of antioxidant protection (Averill-Bates, 2023).

## **2. Challenges and Future Perspectives**

While the potential of secondary metabolites as antioxidants in medicine and agriculture is exciting, several challenges need to be addressed:

**2.1. Bioavailability:** The bioavailability of secondary metabolites can be limited, making it necessary to develop formulations that improve their absorption and effectiveness.

**2.2. Standardization:** Ensuring consistent quality and content of secondary metabolites in herbal medicines and dietary supplements is essential for their therapeutic efficacy and safety.

**2.3. Environmental Concerns:** The cultivation and extraction of secondary metabolite-rich plants can have environmental impacts. Sustainable practices and alternative sources need to be explored.

**2.4. Regulatory Issues:** The regulatory framework surrounding the use of secondary metabolites in medicine and agriculture is complex and varies by region. Harmonization and clear guidelines are essential.

**2.5. Resistance:** Pests and pathogens can develop resistance to secondary metabolite-based pesticides. Integrated pest management strategies are necessary to address this challenge.

### 3. Conclusion

Oxidative stress, a common denominator in various diseases, presents a complex challenge in modern healthcare. Phytomedicine, with its rich repertoire of antioxidant phytochemicals, has emerged as a promising strategy to counteract oxidative stress-induced damage. The mechanisms of action of phytochemicals are diverse, encompassing direct ROS scavenging, enhancement of endogenous antioxidant defenses, and modulation of inflammation. Phytomedicine interventions show potential in preventing and managing oxidative stress-related conditions, including cardiovascular diseases, neurodegenerative disorders, cancer, and metabolic syndrome.

The future of phytomedicine research holds exciting prospects, including personalized nutrition approaches, innovative drug delivery systems, and combination therapies that leverage the synergistic effects of various phytochemicals. As our understanding of the complex interplay between oxidative stress and health deepens, phytomedicine is poised to play an increasingly vital role in the prevention and treatment of diseases associated with oxidative stress.

### 4. Conflict of Interest

The authors report no conflicts of interest.

### 5. References

1. Badyal S., Singh H., Yadav A.K., Sharma S., Bhushan I., 2020. Plant secondary metabolites and their uses. *Plant Archives*, 20:3336-3340.
2. Maleki M., Ghorbanpour M., Kariman K., 2017. Physiological and antioxidative responses of medicinal plants exposed to heavy metals stress. *Plant Gene*, 11:247-254.
3. Leenders F., Groen N., de Graaf N., Engelse M.A., Rabelink T.J., de Koning E.J., Carlotti F., 2021. Oxidative stress leads to  $\beta$ -cell dysfunction through loss of  $\beta$ -cell identity. *Front in Immunology*, 12:690379.
4. Martins S.G., Zilhao R., Thorsteinsdottir S., Carlos A.R., 2021. Linking oxidative stress and DNA damage to changes in the expression of extracellular matrix components. *Frontiers*, 12.
5. McCord J.M., 2000. The evolution of free radicals and oxidative stress. *The American Journal of Medicine*, 108(8): 652-659.
6. Bhatti J.S., Bhatti G.K., Reddy P.H., 2017. Mitochondrial dysfunction and oxidative stress in metabolic disorders- A step towards mitochondria based therapeutic strategies. *Biochimica et Biophysica Acta (BBA)- Molecular Basis of Disease*, 1863(5): 1066-1077.
7. Whitehead N.P., Yeung E.W., Froehner S.C., Allen D.G., 2010. Skeletal muscle NADPH oxidase is increased and triggers stretch-induced damage in the mdx mouse. *PLoS One*, 5:e15354.
8. Mohamed J.B., Hunter M.I.S., 1986. Plasma antioxidants and lipid peroxidation products in duchenne muscular dystrophy. *Clinica Chimica Acta*, 155(2): 123-131.
9. Kowalczyk P., Sulejczak D., Kleczkowska P., Bukowska-Ośko I., Kucia M., Popiel M., Wietrak E., Kramkowski K., Wrzosek K., Kaczyńska K., 2021. Mitochondrial oxidative stress- A causative factor and therapeutic target in many diseases. *International Journal of Molecular Sciences*, 22(24): 13384.
10. Pizzino G., Irrera N., Cucinotta M., Pallio G., Mannino F., Arcoraci V., Squadrito F., Altavilla D., Bitto A., 2017. Oxidative stress: Harms and benefits for human health. *Oxidative Medicine and Cellular Longevity*, 8416763.
11. Ryter S.W., Tyrrell R.M., 1998. Singlet molecular oxygen ( $^1O_2$ ): A possible effector of eukaryotic gene expression. *Free Radical Biology and Medicine*, 24(9): 1520-1534.
12. Gammone M.A., Riccioni G., D'Orazio N., 2015. Marine carotenoids against oxidative stress: effects on human health. *Marine drugs*, 13(10): 6226-46.
13. Özcan P., Fıçıcıoğlu C., Yıldırım O.K., Özkan F., Akkaya H., Aslan I., 2015. Protective effect of resveratrol against oxidative damage to ovarian reserve in female Sprague-Dawley rats. *Reproductive Biomedicine Online*, 31(3): 404-10.

14. Puthur J.T., Sameena P.P., Anjitha K.S., 2021. Functional aspects of plant secondary metabolites in metal stress tolerance and their importance in pharmacology. *Plant Stress*, 2:100038.
15. Gorni P.H., Pacheco A.C., Moro A.L., Silva J.F.A., Moreli R.R., de Miranda G.R., Pelegrini J.M., Zaniboni C.B., Spera K.D., Junior J.L.B., da Silva R.M.G., 2020. Elicitation improves the leaf area, enzymatic activities, antioxidant activity and content of secondary metabolites in *Achillea millefolium* L. grown in the field. *Journal of Plant Growth Regulation*, 40:1652-1666.
16. Bhattacharjee S., Banik N., 2020. Complementation of ROS scavenging secondary metabolites with enzymatic antioxidant defense system augments redox-regulation property under salinity stress in rice. *Physiology and Molecular Biology of Plants*, 26(8):1623-1633.
17. Averill-Bates D.A., 2023. The antioxidant glutathione. *Vitamins and hormones*, 121:109-141.